

# ANALYSIS QUALIFYING EXAM

JANUARY 19, 2012

## REAL ANALYSIS

Answer all 4 questions. In your proofs, you may use any major theorem, except the fact you are trying to prove (or a variant of it). State clearly what theorems you use. Good luck.

### Question 1

- a) Let  $f \in L^1(\mathbb{R})$  and  $F(x) = \int_1^x f(t) dt$ . Then  $F$  is continuous.
- b) Show for  $a > 0$   $\int_1^{\infty} e^{-x^2} \cos(ax) dx = \frac{1}{2} e^{-a^2/4}$ .

### Question 2

State and prove the Hahn-Decomposition Theorem for signed measures.

### Question 3

Let  $1 < p < \infty$  and  $m$  be Lebesgue measure. Let  $M$  be a closed subspace of  $L^p([0; 1]; m)$  such that  $M$  is contained in the space of continuous functions  $C([0; 1])$  (i.e. every element of  $M$  has a continuous function in its equivalence class, which is necessarily unique). Show that there exists  $\epsilon_p > 0$  such that for all  $f \in M$

$$\|f\|_{\infty} \leq C_p \|f\|_p$$

where  $\| \cdot \|_{\infty}$  is the sup norm on  $C([0; 1])$  and  $\| \cdot \|_p$  is the  $L^p$  norm on  $L^p([0; 1]; m)$ .

### Question 4

- a) Prove the uniqueness of a (left-invariant) Haar measure on a locally compact Hausdorff topological group.
- b) Prove that Haar measure for a compact group or abelian group is both left and right invariant.